ATTACK!

Controlice®
HEAD RINSE

SCHEDULING STATUS
To be allocated.

PROPRIETARY NAME AND DOSAGE FORM
Controlice® Head Rinse (Solution).

COMPOSITION
Each 100 ml contains: Dimethicone 4 % (m/m).
Inactive Ingredients: Cyclopentasiloxane and isopropyl myristate.

PHARMACOLOGICAL CLASSIFICATION
D.32.7 Other substances or agents - Topical applications of delousing agents.
Health Supplement.

INDICATION
Controlice® Head Rinse contains the active ingredient, dimethicone which can assist with the topical management of head lice (*Pediculus capitis*) infestation. Efficacy of support may vary between users.

CONTRAINDICATIONS
Do not use if known hypersensitivity or allergy exists towards any of the ingredients.

WARNINGS AND SPECIAL PRECAUTIONS
FOR EXTERNAL USE ONLY. Do not use if known hypersensitivity or allergy exists towards any of the ingredients. Do not apply to damaged or broken skin. Avoid contact with eyes and mucous membranes. If product gets into eyes, wash out immediately with water, and seek medical advice. Upon accidental ingestion, do not induce vomiting. Drink plenty of water, and seek medical advice. Use only as directed. Do not use after date of expiry. See expiry date on unit. Keep away from open flames.
Porphyria: Safety has not been established.
STORE OUT OF REACH OF CHILDREN.

INTERACTIONS
None reported.

PREGNANCY AND LACTATION
Safety for use during pregnancy and lactation (breastfeeding) has not been established.

DOSAGE AND DIRECTIONS FOR USE
FOR EXTERNAL USE ONLY.
Use only as directed.
Keep away from open flames.
Adults and children 1 year and older:
1. Test for sensitivity on the inside of the forearm.
2. Apply sufficient solution to dry hair and spread evenly from the roots to the tips.
3. Ensure scalp and hair are fully covered.
4. Comb hair thoroughly with a Controlice® Comb, area by area.
5. Comb from the base of the hair, as near to the scalp as possible, through to the hair tip to remove all nits (eggs) and dead head lice.
6. Allow hair to dry naturally. Leave on the hair for at least 8 hours or overnight while sleeping.
7. Thereafter wash hair with Controlice® Defence Shampoo or normal shampoo.
8. **Important:** Repeat after 7 days to ensure that the full life cycle of the lice are covered.

SIDE EFFECTS
If you experience any side effects or sensitivity towards any of the ingredients, discontinue use. If symptoms persist, or if any adverse reactions occur, consult your medical practitioner. Side effects include, but are not limited to allergic skin reactions.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT
There are no known symptoms of overdosage. However, if irritation or any other adverse reactions occur, please consult a medical practitioner. Treatment is symptomatic and supportive.

IDENTIFICATION
Clear, watery solution.

PRESENTATION
100 ml solution in a plastic bottle with a spout, in a unit carton, with a package insert.

STORAGE INSTRUCTIONS
Store below 25 °C. Store away from direct sunlight.
Keep away from open flames.
STORE OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER
To be allocated.

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION
Nativa (Pty) Ltd
260 Cradock Avenue, Lyttelton, Centurion, 0157, Gauteng, South Africa
Tel: + 27 (0) 12 664 7110 • Fax: + 27 (0) 12 664 8031
Customer Care Line: **0860 NATIVA (628 482)**
E-mail: health@nativa.co.za • Website: www.nativa.co.za • www.controlice.co.za

DATE OF PUBLICATION OF THE PACKAGE INSERT
August 2007

This medicine has not been evaluated by the Medicines Control Council. This medicine is not intended to diagnose, treat, cure or prevent any disease.

PPIL031/03

Send Head Lice packing:

ATTACK with – Controlice® Oil Spray/Controlice® Head Rinse
COMB with – Controlice® Lice Comb/Controlice® Lice Buster
PROTECT with – Controlice® Defence Shampoo/ Controlice® Defence Spray

LICE FACTS

Head lice are small, fast crawling parasites that feed on human blood. They're clear in colour when first hatched, and then quickly develop a reddish-brown colour after feeding. An adult louse is about the size of a pin head and their nits (eggs) are about two thirds smaller. In order to survive, lice need to feed on the blood of their hosts every 3-4 hours. While on their hosts they also extend the lice population by mating every 3-4 hours. On average they stay alive between 23-30 days.

WHO GETS HEAD LICE AND HOW DO THEY GET THEM?

Only humans can get head lice. If stranded away from the human scalp they die quickly (1-2 days). The usual way people are infested is from direct hair-to-hair contact with another person. Head lice don't have wings therefore they cannot fly or jump from head to head without assistance, but they are terrific runners and they can hide. Head lice have six legs and grasp the hair with claws.

Lice infestations can come from almost anywhere:
A classmate, another child's clothing, hat, comb or brush.
Lice outbreaks are common among school children and even the cleanest child can easily become infested. During a lice infestation, it is important that you INSPECT all family members for at least two weeks.

Head lice should be suspected when there is intense itching of the scalp and/or at the nape of the neck. You will see small, whitish nits attached to the hair shaft. Head lice may also be indicated if one finds tiny, black spots on bed pillows, sheets or clothing near the neckline and shoulders. These black spots consist of digested blood excreted by a head louse after feeding.

ATTACK:
The active ingredients in Controlice® Oil Spray / Controlice® Head Rinse can be used to eliminate head lice already infesting the hair. It is very important to use the product as directed. After using the Controlice Oil® Spray / Controlice® Head Rinse, it is necessary to comb out all the nits and dead head lice with a Controlice® Comb. The Controlice® Combs are precision engineered products that are very effective in removing nits. The unique tooth configuration provides a gentle scraping action which dislodges lice and nits without damaging the hair. Use the Controlice® Lice Buster for long hair and the Controlice® Lice Comb for short hair.

Important: Repeat treatment and combing after 7 days.

PROTECT:
Controlice® Defence Shampoo and / or Controlice® Defence Spray contain active ingredients which repel head lice and protect against head lice infestation during a lice outbreak. It can be used by adults and children. Use the Controlice® Combs on a regular basis to check for head lice.





