Proprietary Name and Dosage Form:
OsteoEze Acute (Tablets)

Composition

Active Ingredients	Per tablet:	Per 3 tablets (maximum daily dosage):
Glucosamine Sulfate KCl	400 mg	1200 mg
Boswellia serrata Extract	100 mg	300 mg
Curcuma longa (Curcumin 95 %) Root Extract	100 mg	300 mg
Zingiber officinale (Ginger) Root Extract (4:1)	100 mg	300 mg

Pharmacological Classification
Combination Product Western Herbals.

Indications
OsteoEze® Acute is a general supplement for joints that contains glucosamine, *Boswellia serrata*, curcumin and ginger. This combination of active ingredients can assist the joints with acute pain management and mobility support. In addition, it has anti-inflammatory properties that can assist with swelling reduction. Efficacy of support may vary between users.

Contraindications
Do not use this product if allergic to shellfish. Do not use if known hypersensitivity or allergy exists towards any of the ingredients.

Warnings
Do not use this product if allergic to shellfish. Do not use if known hypersensitivity or allergy exists towards any of the ingredients. People using anticoagulant - or antiplatelet medication, such as Warfarin, should use this product with caution as curcumin and glucosamine may have a blood thinning effect. Do not exceed the recommended daily dosage. Do not use after date of expiry. Products containing herbal ingredients may interact with some medication. If you are taking any chronic medication, do not use this product without consulting a medical practitioner.
Porphyria: Safety has not been established.
KEEP OUT OF REACH OF CHILDREN.

Interactions
People using anticoagulant - or antiplatelet medication, such as Warfarin, should use this product with caution as *Curcuma longa* (Curcumin 95 %) Root Extract and glucosamine may have a blood thinning effect.

Pregnancy and Lactation
Safety for use during pregnancy and lactation (breastfeeding) has not been established.

Dosage and Directions for Use
Do not exceed the recommended daily dosage.
Adults 18 years and older:
Take two tablets three times a day after meals, with a glass of water, for two days. Thereafter, take one tablet three times a day after meals, with a glass of water.

Side Effects and Special Precautions
If you experience any side effects or sensitivity towards any of the ingredients, discontinue use. If symptoms persist, or if any adverse reactions occur, consult a medical practitioner. Side effects may include, but are not limited to, nausea, vomiting, diarrhoea, constipation and dizziness.

Known Symptoms of Overdosage and Particulars of Its Treatment
Do not exceed the recommended daily dosage. If an accidental overdose occurs, please consult a medical practitioner. Treatment is symptomatic and supportive.

Identification
Oblong-shaped, film-coated, blue tablet.

Presentation
Blister pack with 3 strips of 10 tablets each, in a unit carton

Storage Instructions
Store below 25 °C. Store away from moisture and direct sunlight.
KEEP OUT OF REACH OF CHILDREN.

Complementary Registry Number
005101

Name and Business Address of Applicant
Nativa (Pty) Ltd
Private Bag X 1030, Lyttelton, 0140, Gauteng, South Africa
Tel: +27 (0) 12 664 7110 • Fax: +27 (0) 12 664 8031
Customer Care Line: **0860 NATIVA (628 482)**
E-mail: health@nativa.co.za
Website: www.nativa.co.za

Date of Publication of the Package Insert
April 2006

This medicine has not been evaluated by the Medicines Control Council.
This medicine is not intended to diagnose, treat, cure or prevent any disease.